arXiv:hep-ph/0211400v1 26 Nov 2002

Single Transverse Spin Asymmetry in $p^{\uparrow}p \to \pi X$ and $ep^{\uparrow} \to \pi X$

Yuji Koike[a]MCSD]Department of Physics, Niigata University, Niigata 950-2181, Japan

[a][

Single transverse spin asymmetry for the pion production in pp and ep collisions is studied in the framework of collinear factorization. The soft-gluon pole contributions with twist-3 distribution and fragmentation functions are identified and its characteristic features are discussed.

In this report we discuss the single transverse-spin asymmetry A_N for the pion production with large transverse momentum in $pp^{\uparrow} \to \pi(\ell)X$ and $p^{\uparrow}e \to \pi(\ell)X$ relevant for RHIC-SPIN, HERMES and COMPASS experiments *etc.* According to the QCD factorization theorem, the polarized cross section for $pp^{\uparrow} \to \pi X$ consists of three twist-3 contributions:

$$\begin{array}{ll} (A) & G_a(x_1,x_2) \otimes q_b(x') \otimes D_c(z) \otimes \hat{\sigma}_{ab\to c}, \\ (B) & \delta q_a(x) \otimes E_b(x_1',x_2') \otimes D_c(z) \otimes \hat{\sigma}'_{ab\to c}, \\ (C) & \delta q_a(x) \otimes q_b(x') \otimes \widehat{E}_c(z_1,z_2) \otimes \hat{\sigma}''_{ab\to c}, \end{array}$$

where the functions $G_a(x_1,x_2)$, $E_b(x_1',x_2')$ and $\widehat{E}_c(z_1,z_2)$ are the twist-3 quantities representing, respectively, the transversely polarized distribution, the unpolarized distribution, and the fragmentation function for the pion. Other functions are twist-2: $q_b(x')$ is the unpolarized quark or gluon distribution, $\delta q_a(x)$ is the transversity distribution in $p^{\uparrow}$, and $D_c(z)$ is the fragmentation function for the pion. a, b and c stand for the parton's species, sum over which is implied. δq_a, E_b and $\widehat{E}_c$ are chiral-odd. $\hat{\sigma}_{ab\to c}$ *etc* are the partonic hard cross section producing large transverse momentum of the pion. Corresponding to the above (A) and (C), the polarized cross section for $ep^{\uparrow} \to \pi X$ (final electron is not detected) receives two twist-3 contributions:

$$\begin{array}{ll} (A') & G_a(x_1,x_2) \otimes D_a(z) \otimes \hat{\sigma}_{ea\to a}, \\ (C') & \delta q_a(x) \otimes \widehat{E}_a(z_1,z_2) \otimes \hat{\sigma}'_{ea\to a}. \end{array}$$

The (A) and (B) contributions for $pp^{\uparrow} \to \pi X$ have been analyzed in [1] and [2], respectively, and it has been shown that (A) gives rise to large A_N at large x_F as observed in E704, and (B) is negligible in all kinematic region. Here we extend the study to the (C) term at RHIC energy and also the asymmetry in ep collision.

The transversely polarized twist-3 distribution $G_F^a(x_1,x_2)$ relevant to G_a in the (A) and (A') terms is given in [2], and the twist-3 fragmentation function $\widehat{E}_F^c(z_1,z_2)$ for $\widehat{E}_c$ in (C) and (C') is defined in [4]. According to a detailed analysis, G_F^a and $\widehat{E}_F^c$ contribute to the cross section at $x_1 = x_2 = x$ and $z_1 = z_2 = z$, respectively (soft-gluon-pole contribution). In particular, the derivarives $(d/dx)G_F^a(x,x)$ and $(d/dz)\widehat{E}_F^c(z,z)$ appear in the cross section formula. In the large x_F region, i.e. for the production of pion in the forward direction of the polarized nucleon, the main contribution comes from the large-x and large-z region of distribution and fragmentation functions, respectively. Since G_F^a and $\widehat{E}_F^c$ behaves as $G_F^a(x,x) \sim (1-x)^\beta$ and $\widehat{E}_F^c(z,z) \sim (1-z)^{\beta'}$ with β, $\beta' > 0$, $|(d/dx)G_F^a(x,x)| \gg |G_F^a(x,x)|$, $|(d/dz)\widehat{E}_F^c(z,z)| \gg |\widehat{E}_F^c(z,z)|$ at large x and z. In particular, the valence component of G_F^a and $\widehat{E}_F^c$ dominate in this region. We thus keep only the valence quark contribution for the derivative of these soft-gluon pole functions ("valence-quark soft-gluon approximation") for the pp collision. For the ep case, we include all the soft-gluon pole contribution.

In general A_N is a function of $S = (P+P')^2 \simeq 2P\cdot P'$, $T = (P-\ell)^2 \simeq -2P\cdot\ell$ and $U = (P'-\ell)^2 \simeq -2P'\cdot\ell$ where P, P' and ℓ are the momenta of $p^\uparrow$, unpolarized p (or e), and the pion respectively. In the following we use S, $x_F = \frac{2\ell_\parallel}{\sqrt{S}} = \frac{T-U}{S}$ and $x_T = \frac{2\ell_T}{\sqrt{S}}$ as independent variables. The cross section formula for the (C) term is given in [4].



Figure 1. A_N^{pp} at $\sqrt{S} = 20$ GeV and $\ell_T = 1.5$ GeV together with E704 data.

To estimate the (C) contribution, we introduce a simple model ansatz as $\widehat{E}_F^a(z,z) = K_a D_a(z)$ with a flavor dependent factor K_a. Assuming that the (C) is the sole origin of A_N, K_a's are determined to be $K_u = -0.11$ and $K_d = -0.19$ so that it can approximately reproduce A_N^{pp} observed in E704 data at $\sqrt{S} = 20$ GeV and $\ell_T = 1.5$ GeV. We refer the readers to [2] for the adopted distribution and fragmentation functions. The result for A_N^{pp} from the (C) (chiral-odd) term is shown in Fig. 1 in comparison with the (A) (chiral-even) contribution. The magnitude of the (A) term in Fig. 1 is also determined with the ansatz $G_F^a(x,x) = K'_a q_a(x)$ ($K'_u = -K'_d = 0.07$) so that it can approximately reproduce E704 data by itself. Recall that our valence-quark soft-gluon approximation is valid at large x_F. Both effects give rise to A_N^{pp} similar to the E704 data. The origin of the growing A_N at large x_F is (i) large partonic cross sections ($\sim 1/\hat{t}^2$ term) and (ii) the derivative of the soft-gluon pole functions.

With the parameters K_a and K'_a fixed, A_N^{pp} at RHIC energy ($\sqrt{S} = 200$ GeV) is shown in Fig.2 at $l_T = 1.5$ GeV. Both (A) and (C) contributions give slightly smaller A_N^{pp} than the STAR data reported at SPIN2002. Fig. 3 shows the ℓ_T-dependence of A_N^{pp} at $\sqrt{S} = 200$ GeV and $x_F = 0.6$, indicating quite large l_T dependence in both (A) and (C) contributions at $1 < l_T < 4$ GeV region, which is typical to the twist-3 effect ($\sim O(1/\ell_T)$). Polarized cross section receives $O(\ell_T/S)$ contribution as well, so that the A_N tends to become more stable at larger ℓ_T.



Figure 2. A_N^{pp} at $\sqrt{S} = 200$ GeV and $\ell_T = 1.5$ GeV.



Figure 3. ℓ_T dependence of A_N^{pp} at $\sqrt{S} = 200$ GeV and $x_F = 0.6$.

We next discuss the asymmetry A_N^{ep} for $p^\uparrow e \to \pi(\ell)X$ where the final electron is not observed. In our $O(\alpha_s^0)$ calculation, the exchanged photon remains highly virtual as far as the observed π has a large transverse momentum ℓ_T with respect to the ep axis. Therefore experimentally one needs to integrates only over those virtual photon events ($Q^2 > (S/2)[1 - x_F/\sqrt{x_F^2 + x_T^2}]$) to compare with our formula.



Figure 4. A_N^{ep} at $\sqrt{S} = 20$ GeV and $\ell_T = 1.5$ GeV.

Using the twist-3 distribution and fragmentation functions used to describe pp data, we show in Fig. 4 A_N^{ep} corresponding to (A')(chiral-even) and (C')(chiral-odd) contributions. Remarkable feature of Fig. 3 is that in both chiral-even and chiral-odd contributions (i) the sign of A_N^{ep} is opposite to the sign of A_N^{pp} and (ii) the magnitude of A_N^{ep} is much larger than that of A_N^{pp}, in particular, at large x_F, and it even overshoots one. (In our convention, $x_F > 0$ corresponds to the production of π in the forward hemisphere of the initial polarized proton both in $p^\uparrow p$ and $p^\uparrow e$ case.) The origin of these features can be traced back to the color factor in the dominant diagrams for the *twist-3 polarized* cross sections in ep and pp collisions. Of course, the asymmetry can not exceeds one, and thus our model estimate needs to be modified. First, the applied kinematic range of our formula should be reconsidered: A_N at ℓ_T smaller than a few GeV may not be ascribed to the twist-3 effect. Second, our model ansatz of $G_F^a(x,x) \sim q^a(x)$ and $\widehat{E}_F^a(z,z) \sim D^a(z)$

is not appropriate at $x \to 1$ and $z \to 1$, respectively. The derivative of these functions, which is important for the growing A_N^{pp} at large x_F, eventually leads to divergence of A_N^{pp} at $x_F \to 1$ as $\sim 1/(1-x_F)$.

As a possible remedy to this pathology we tried the following. For the (A) and (A') (chiral-even) contribution, as an example, we have a model $G_F^a(x,x) \sim q_a(x) \sim_{x\to 1} (1-x)^{\beta_a}$ where $\beta_u = 3.027$ and $\beta_d = 3.774$ in the GRV distribution we adopted. Tentatively we shifted $\beta_{u,d}$ as $\beta_a \to \beta_a(x) = \beta_a + x^3$, which suppresses the divergence of A_N at $x_F \to 1$ but still causes rising behavior of A_N at large x_F. The result for A_N with this modification of $G_F(x,x)$ is shown in Fig. 5 (A_N^{ep}), Fig. 6 (A_N^{pp} at E704 energy) and Fig. 7 (A_N^{pp} at RHIC energy). This avoids overshooting of one in A_N^{ep} but reduces A_N^{pp}, typically by factor 2, which makes the deviation from E704 and STAR data larger. A similar modification of $\widehat{E}_F(z,z)$ in the (C) and (C') (chiral-odd) contribution also leads to the same change in A_N.



Figure 5. A_N^{ep} with a modified G_F.



Figure 6. A_N^{pp} with a modified G_F at $\sqrt{S} = 20$ GeV.



Figure 7. A_N^{pp} with a modified G_F at $\sqrt{S} = 200$ GeV.

To summarize we have studied the A_N for pion production in *pp* and *ep* collisions. We have pointed out that both (A)(chiral-even) and (C)(chiral-odd) ((A') and (C') for *ep* collisions) contributions can be equally important sources for the asymmetry. Although our approach provides a systematic framework for the large ℓ_T production, applicability of the formula to the currently available low ℓ_T data still needs to be tested, in particular, detailed comparison of ℓ_T-dependence and the global analysis of *pp* and *ep* data is necessary.

Acknowledgement: This work is supported in part by the Grant-in-Aid for Scientific Research of Monbu-Kagaku-sho.

REFERENCES

1. J. Qiu and G. Sterman, Phys. Rev. **D59** (1999) 014004.
2. Y. Kanazawa and Y. Koike, Phys. Lett. **B478** (2000) 121; **B490** (2000) 99.
3. Y. Koike, hep-ph/0106260 (Proceedings of DIS2001, Bologna, Italy, April, 2001.)
4. Y. Koike, hep-ph/0210396 (Proceedings of SPIN2002, Long Island, USA, Sep. 2002)